

Critical Properties of One-Dimensional Electron Systems with $1/r^\alpha$ -type Long-Range Interactions

Yasumasa TSUKAMOTO and Norio KAWAKAMI

Department of Applied Physics, Osaka University, Suita, Osaka 565-0871, Japan

(Received)

We study the critical properties of one-dimensional electron systems with the $1/r^\alpha$ -type long-range interaction ($\alpha \geq 1$). Using bosonization methods, we discuss how the 1D system exhibits a crossover from the Tomonaga-Luttinger liquid to the Wigner crystal when $\alpha \rightarrow 1$. In particular, a detailed analysis is given to the Fermi-edge singularity phenomena when a *mobile* core hole is created suddenly in the valence band. It is found that the effect due to the core-hole motion on the Fermi-edge singularity is gradually suppressed as the system approaches the Wigner crystal, and its critical behavior is dominated by the ordinary impurity forward-scattering inherent in a static hole.

KEYWORDS: long-range interaction, bosonization, Fermi-edge singularity

§1. Introduction

Since the recent advances in technology made it possible to fabricate quantum wires, the study on one-dimensional (1D) electron systems has been one of the most attractive research areas in condensed matter physics. Low-energy properties of 1D correlated electron systems may be generally described by Tomonaga-Luttinger (TL) liquid theory. [1,2,3,4] In this framework, the interaction among electrons is assumed to be short-ranged, which may result from the cooperative screening effect. However, this assumption is not always valid for quasi-1D systems because the screening effect due to electrons becomes poorer in low-dimensional systems, and thereby the long-range nature of the interaction may manifest itself in various quantities. In fact, the optical experiments for quantum wires [5] suggest that the effect of long-range interaction originating from the unscreened Coulomb force should be taken into account in treating 1D systems theoretically. More recently, it has been recognized that the unscreened long-range interaction is also important for quasi-1D Carbon nanotubes. [6,7,8]

Stimulated by these facts, many theoretical works on 1D electron systems with the $1/r$ Coulomb interaction have been reported. Among others, Schulz [9] has clarified that the $1/r$ long-range interaction enhances the $4k_F$ charge density correlation, driving the system to the *Wigner crystal* which shows a critical behavior quite different from the ordinary TL liquid. Subsequently, various properties in 1D systems with $1/r$ interaction have been examined. [10,11,12,13,14] For instance, [10] a simple power-law temperature dependence in the conductance G for the TL liquid is changed to $G \sim \exp[-\ln^{3/2}(1/T)]$, which vanishes with temperature faster than any power law, for the Wigner crystal

with the $1/r$ interaction.

In this paper, we investigate the critical behavior of 1D electron systems with the long-range interaction by means of abelian bosonization methods. To clearly see how the TL liquid is changed to the Wigner crystal, we systematically deal with the $1/r^\alpha$ -type long-range interaction with a continuously changing exponent α . In particular, special attention is focused on the Fermi-edge singularity phenomena [13] with a *mobile* core hole. We discuss the effect of the core-hole motion on the singularity by computing the correlation functions for the orthogonality catastrophe (OC) and the Fermi-edge singularity (FES). It is found that the effect of a mobile core hole is *suppressed* in the 1D system with long-range interaction, which is contrasted to our previous finding [15] that the effect of the core-hole motion plays an important role in TL liquids with short-range interaction.

The paper is organized as follows. In §2, we introduce the model with the $1/r^\alpha$ -type interaction, which also contains the impurity scattering due to a mobile core hole. Then in §3, we clarify how the mobile core hole created affects the long-time critical properties in the OC and the FES, by observing a crossover behavior from the TL liquid to the Wigner crystal. A brief summary of the results is given in §4.

§2. Model with $1/r^\alpha$ Long-range Interaction

We consider a 1D correlated electron system with $1/r^\alpha$ -type long-range interaction. As mentioned above, the screening effect is not complete in quasi-1D systems, so that the bare interaction is modified from its pure Coulombic form in a non-trivial way. For instance, it has been suggested that for large distance the interaction might show the power-law behavior. [16] We shall consider here the $1/r^\alpha$ interaction with $\alpha \geq 1$, which can

systematically describe a crossover from the TL liquid to the Wigner crystal. Since the $1/r^\alpha$ interaction exhibits a singular property for $r \rightarrow 0$, we introduce the interaction with short-range cut off d , [9]

$$V(r) = \frac{1}{(r^2 + d^2)^{\nu+1/2}}, \quad \nu \geq 0, \quad (2.1)$$

which is Fourier transformed as,

$$V(p) = \frac{2\sqrt{\pi}}{\Gamma(\nu + 1/2)} \left| \frac{p}{2d} \right|^\nu K_\nu(pd), \quad (2.2)$$

where Γ is the gamma function and K_ν denotes the ν -th order modified Bessel function. Here we have introduced the index ν ($\alpha = 2\nu + 1$), which is convenient for the following discussions. By continuously changing ν , we can alter the range of the interaction; For instance, in the $\nu \rightarrow \infty$ limit we can effectively describe the δ -function type interaction. For $\nu > 0$ case, the uniform component of the interaction is given by,

$$V(0) = \frac{\sqrt{\pi}\Gamma(\nu)}{2d^{2\nu}\Gamma(\nu + 1/2)} \equiv F(\nu, d) \quad (2.3)$$

which is the key quantity to control various correlation exponents so far as the system belongs to the TL liquid.

The low-energy effective Hamiltonian for the host part is now written down in the g -ology description, [1, 2, 3, 4]

$$\begin{aligned} \mathcal{H}_{total} &= \mathcal{H}_\rho + \mathcal{H}_\sigma \\ \mathcal{H}_\rho &= \mathcal{H}_{free}^\rho + \mathcal{H}_{g_2}^\rho + \mathcal{H}_{g_4}^\rho, \end{aligned}$$

which consists of the charge (ρ) and spin (σ) sectors. The free part of the charge sector has the ordinary harmonic form with respect to the charge density $\rho_+(p)$ ($\rho_-(p)$) for the right- (left-) going electrons,

$$\begin{aligned} \mathcal{H}_{free}^\rho &= \frac{\pi v_F}{L} \sum_{p \neq 0} \sum_{r=\pm} : \rho_r(p) \rho_r(-p) : \\ &+ \frac{\pi}{2L} [v_{N_\rho} (N_{+\rho} + N_{-\rho})^2 + v_{J_\rho} (N_{+\rho} - N_{-\rho})^2] \end{aligned} \quad (2.4)$$

where v_F is the Fermi velocity and $v_{N_\rho} = v_{J_\rho} = v_F$. The long-range interaction defined by Eq.(2.2) leads to the forward scattering terms for the charge part,

$$\mathcal{H}_{g_2}^\rho = \frac{2}{L} \sum_p g_2(p) \rho_+(p) \rho_-(-p) \quad (2.5)$$

$$\mathcal{H}_{g_4}^\rho = \frac{1}{L} \sum_{p, r=\pm} g_4(p) : \rho_r(p) \rho_r(-p) : \quad (2.6)$$

where $g_2(p)/\pi v_F = g_4(p)/\pi v_F = V(p)$. Here we have neglected the backward and Umklapp scatterings, since they are not relevant in the present case. The spin part is given by SU(2) critical Hamiltonian, [2, 3, 4] the explicit form of which is not necessary in the following treatment.

Let us now introduce the effective impurity scattering which describes the interaction between 1D electrons and a *mobile* core-hole created suddenly, e.g. by the X-ray absorption. In the previous paper, [15] we have derived the following effective Hamiltonian for the impurity scattering,

tering,

$$\begin{aligned} \mathcal{H}_I^\rho &= \frac{v_f^{(1)}}{L} \sum_p \tilde{V}(p) [\rho_+(p) + \rho_-(p)] \\ &+ \frac{v_f^{(2)}}{L} \sum_p [\rho_+(p) - \rho_-(p)]. \end{aligned} \quad (2.7)$$

The first term is the ordinary forward scattering between electrons and an impurity, for which we may change the spatial dependence of the impurity potential by tuning the momentum dependent part $\tilde{V}(p)$. The second non-trivial term does not exist in the microscopic model, but is induced by the motion of the core hole; namely, it originates from the coupling between electron currents and the core-hole motion. [15] Thus the interaction strength $v_f^{(2)}$ is dependent on the momentum of the mobile core hole. For instance, if the impurity is assumed to be static(localized), then $v_f^{(2)} = 0$. This term, which is referred to as the *asymmetric* forward scattering, plays an important role for FES in TL liquids. It was found [15] that the asymmetric scattering is to be short-ranged, even if the microscopic interaction is long ranged. In order to distinguish $v_f^{(1)}$ with $v_f^{(2)}$, we henceforth refer to the former as the *symmetric* forward scattering.

Following standard procedures in the bosonization, [4] $\mathcal{H}_\rho$ can be easily diagonalized. Introducing the unitary operator U ($U\mathcal{H}_\rho U^\dagger \equiv \mathcal{H}_\rho^D$), we have

$$\begin{aligned} \mathcal{H}_\rho^D &= \frac{\pi}{L} \sum_{p \neq 0} \sum_{r=\pm} v_\rho(p) : \rho_r(p) \rho_r(-p) : \\ &+ \frac{\pi}{2L} [v'_{N_\rho} (N_{+\rho} + N_{-\rho})^2 + v'_{J_\rho} (N_{+\rho} - N_{-\rho})^2] \end{aligned} \quad (2.8)$$

where we define $v'_{N_\rho} = v_\rho/K_\rho$ and $v'_{J_\rho} = v_\rho K_\rho$ in terms of the TL parameter K_ρ and the charge velocity v_ρ as,

$$K_\rho(p) = \frac{1}{\sqrt{1 + 2V(p)}}, \quad v_\rho(p) = v_F \sqrt{1 + 2V(p)}. \quad (2.9)$$

The impurity potential $\mathcal{H}_I^\rho$ is simultaneously transformed as,

$$\begin{aligned} \mathcal{H}_\rho^{eff} &\equiv U\mathcal{H}_I^\rho U^\dagger \\ &= \frac{v_f^{(1)}}{L} \sum_p \tilde{V}(p) \sqrt{K_\rho(p)} [\rho_+(p) + \rho_-(p)] \\ &+ \frac{v_f^{(2)}}{L} \sum_p \frac{1}{\sqrt{K_\rho(p)}} [\rho_+(p) - \rho_-(p)]. \end{aligned} \quad (2.10)$$

Notice that the p dependence of K_ρ dramatically modifies the form of correlation functions. [9, 13]

§3. Correlation Functions with a Core Hole Suddenly Created

Let us now discuss the critical properties for 1D systems with $1/r^\alpha$ interaction. We here focus on the FES phenomena with a *mobile* core hole which is created in the valence band by the X-ray absorption. [15, 17] In particular, by analyzing the effects of the symmetric scatter-

ing $v_f^{(1)}$ and asymmetric scattering $v_f^{(2)}$ in Eq.(2.7), we clarify how these terms affect the correlation functions related to the FES. We also observe the crossover to the Wigner crystal by continuously changing the power α in the interaction of $1/r^\alpha$. Note that the FES phenomena for the $1/r$ interaction have been discussed in detail for a static core hole by Otani and Ogawa. [13]

To compute the correlation functions related to the FES, we introduce a core hole created suddenly at $t = 0$, which thereby gives rise to the impurity potential $\mathcal{H}_p^I$. Then the resulting time-dependent correlation functions can be calculated by standard methods [18, 19, 20]. Here we employ the method suitable for our formulation based on the bosonization, i.e. the *boundary condition changing operator* (BCCO) method. [21] The BCCO is a kind of the unitary transformations which is introduced so that the total Hamiltonian with impurity scattering $\mathcal{H}_p^D + \mathcal{H}_p^{eff}$ should be transformed into a free one. According to Affleck and Ludwig, [21] we can write down the BCCO $U_M(t)$ of our system as $U_M(t) = \exp[-iS_M(t)]$ with

$$S_M(t) = \frac{\pi i}{L} \sum_{p \neq 0} \frac{e^{-\alpha|p|/2}}{p} \times [F_+(p)\rho_+(p)e^{ipv_\rho(p)t} - F_-(p)\rho_-(p)e^{-ipv_\rho(p)t}]$$

where

$$F_\pm(p) = \frac{1}{\sqrt{2K_\rho(p)}} \left(\frac{v_f^{(1)}}{\pi} \theta_1(p) \right) \pm \sqrt{2K_\rho(p)} \left(\frac{v_f^{(2)}}{\pi} \theta_2(p) \right).$$

Here $\theta_1(p)$ and $\theta_2(p)$ are given as,

$$\theta_1(p) = \frac{\sqrt{2}K_\rho(p)}{v_\rho(p)} \tilde{V}(p) = \frac{\sqrt{2}\tilde{V}(p)}{1 + 2V(p)} \equiv \Theta(p) \quad (3.2)$$

$$\theta_2(p) = \frac{1}{\sqrt{2}v_\rho(p)K_\rho(p)} = \frac{1}{\sqrt{2}} \equiv \theta, \quad (3.3)$$

where we have used Eq.(2.9) to derive the second formulae. Notice that since the symmetric scattering has p -dependence like $v_f^{(1)}\tilde{V}(p)$, the corresponding $\theta_1(p)$ remains p -dependent, whereas the asymmetric scattering $v_f^{(2)}$ is assumed to be of δ -function type, so that $\theta_2(p)$ is independent of p .

By using the BCCO, we straightforwardly write down the general formulae for the correlation functions of the orthogonality catastrophe (OC) and the Fermi-edge singularity (FES). Making use of $\Theta(p)$ and θ , the correlation function for the OC, $\langle U_M^\dagger(t)U_M(0) \rangle$, is obtained as,

$$\begin{aligned} & Re \langle U_M^\dagger(t)U_M(0) \rangle \\ &= \exp \left[- \left(\frac{v_f^{(1)}}{\pi} \right)^2 \int_0^\infty dp \frac{e^{-\alpha p}}{p} \frac{1}{4K_\rho(p)} \Theta(p)^2 (1 - \cos pv_\rho(p)t) \right] \\ &\times \exp \left[- \left(\frac{v_f^{(2)}}{\pi} \theta \right)^2 \int_0^\infty dp \frac{e^{-\alpha p}}{p} K_\rho(p) (1 - \cos pv_\rho(p)t) \right]. \end{aligned} \quad (3.4)$$

This quantity characterizes how the density of conduction electron responds when a mobile core hole is suddenly created in the valence band. Note that the factor including $v_f^{(2)}$ is inherent in our system with a mobile core hole. Similarly, the correlation function for the FES, for which an electron is added to the conduction band can be derived as,

$$\begin{aligned} & Re \langle U_M^\dagger(t)\psi_r(0,t)\psi_r^\dagger(0,0)U_M(0) \rangle \\ &= \exp \left[- \int_0^\infty dp \frac{e^{-\alpha p}}{p} \frac{1}{4K_\rho(p)} \left(1 + \frac{v_f^{(1)}}{\pi} \Theta(p) \right)^2 (1 - \cos pv_\rho(p)t) \right] \\ &\times \exp \left[- \int_0^\infty dp \frac{e^{-\alpha p}}{p} K_\rho(p) \left(\frac{1}{2} + r \frac{v_f^{(2)}}{\pi} \theta \right)^2 (1 - \cos pv_\rho(p)t) \right] \\ &\times t^{-\frac{1}{2}}, \end{aligned} \quad (3.5)$$

where the last term $t^{-\frac{1}{2}}$ is the contribution from the spin sector with SU(2) symmetry.

Below, we discuss these correlation functions by explicitly evaluating the above integrals for each case, for which a mobile core hole is treated in several circumstances. We first consider the Wigner crystal with $1/r$ interaction as the host system, in which two types of the impurity potential are discussed: (a) $\tilde{V}(p) = V(p)$ and (b) $\tilde{V}(p) = 1$. The former corresponds to the impurity scattering by the $1/r$ potential, while the latter to the δ -function potential. To see the crossover to the Wigner crystal, we then move to the system with the $1/r^\alpha$ interaction ($\nu > 0$) in Eq.(2.1), and calculate the correlation functions for two cases of impurity scattering, (a) $\tilde{V}(p) = V(p)$ case and (b) $\tilde{V}(p) = 1$ cases. Note that the asymmetric forward scattering $v_f^{(2)}$ is always regarded to be short-ranged, as mentioned in §2.

3.1 Wigner crystal: $V(r) = 1/r$ case

Let us begin with the case of the Wigner crystal in which the interaction between host electrons is of the $1/r$ -type ($\nu = 0$ in Eqs.(2.1) and (2.2)).

3.1.1 $1/r$ -type impurity potential

Setting $\tilde{V}(p) = V(p)$, we first regard the impurity scattering due to a mobile core hole to be of the $1/r$ type. Using the relation $V(p) = 2K_0(pd) \simeq 2|\ln|pd|| \gg 1$ for small p , $\Theta(p)$ can be approximated as $\Theta(p) \simeq 1/\sqrt{2}$. Since the TL parameter K_ρ still has the p -dependence, we evaluate Eq.(3.4) by exploiting the generalized formula derived by Otani and Ogawa, [13]

$$\begin{aligned} & Re \langle U_M^\dagger(t)U_M(0) \rangle \\ &\simeq \exp \left[- \frac{1}{3} \left(\frac{v_f^{(1)}}{\sqrt{2}\pi} \right)^2 (\ln(t/d))^{3/2} \right] \\ &\times \exp \left[- \left(\frac{v_f^{(2)}}{\sqrt{2}\pi} \right)^2 (\ln(t/d))^{1/2} \right]. \end{aligned} \quad (3.6)$$

Similarly, we can derive the FES correlation function

as,

$$\begin{aligned} & Re < U_M^\dagger(t) \psi_r(0, t) \psi_r^\dagger(0, 0) U_M(0) > \\ &= \exp \left[-\frac{1}{3} \left(1 + \frac{v_f^{(1)}}{\sqrt{2\pi}} \right)^2 (\ln(t/d))^{3/2} \right] \\ &\times \exp \left[-\left(\frac{1}{2} + r \frac{v_f^{(2)}}{\sqrt{2\pi}} \right)^2 (\ln(t/d))^{1/2} \right] t^{-1/2}. \quad (3.7) \end{aligned}$$

It is seen that the first factor of $\exp[-\text{const.} \ln^{3/2}(t/d)]$ in (3.6) and (3.7), which is caused by the ordinary symmetric scattering $v_f^{(1)}$, gives the most dominant contribution to the correlation functions in the long-time regime. On the other hand, the asymmetric scattering inherent in the mobile core hole results in the subdominant factor, $\exp[-\text{const.} \ln^{1/2}(t/d)]$. This implies that for the Wigner crystal with $1/r$ interaction, the effect of the core-hole motion is substantially suppressed due to the long-range nature of the host interaction. This conclusion is quite contrasted to the ordinary TL liquid case, for which both of the symmetric and asymmetric scatterings equally contribute to the long-time behavior of FES phenomena. [15] We note that our result Eq.(3.7) for a static core hole ($v_f^{(2)} = 0$) agrees with those obtained previously for the $1/r$ case. [13]

3.1.2 δ -function impurity potential

The suppression of the impurity effect occurs more dramatically when we consider the δ -function type impurity potential by setting $\tilde{V}(p) = 1$. Such a problem with δ -function scattering in the Wigner crystal has been treated previously for a static core hole by Fabrizio et al. [10] Though $\Theta(p)$ has still p -dependence in this case, we can calculate the integrals in the same way mentioned above. The OC correlation function takes the asymptotic form,

$$\begin{aligned} & Re < U_M^\dagger(t) U_M(0) > \\ &= \exp \left[\frac{1}{4} \left(\frac{v_f^{(1)}}{\sqrt{2\pi}} \right)^2 (\ln(t/d))^{-1/2} \right] \\ &\times \exp \left[-\left(\frac{v_f^{(2)}}{\sqrt{2\pi}} \right)^2 (\ln(t/d))^{1/2} \right] \dots \quad (3.8) \end{aligned}$$

It should be noted that in contrast to the result obtained in Eq.(3.6), the asymmetric scattering $v_f^{(2)}$ inherent in a mobile core-hole plays the most dominant role for the OC in the long-time regime. However, we can see the dramatic suppression of the mobile impurity effect in the FES correlation function, which is given by,

$$\begin{aligned} & Re < U_M^\dagger(t) \psi_r(0, t) \psi_r^\dagger(0, 0) U_M(0) > \\ &\simeq \exp \left[-\frac{1}{3} (\ln(t/d))^{3/2} \right] \cdot \exp \left[-\left(\frac{v_f^{(1)}}{\sqrt{2\pi}} \right)^2 (\ln(t/d))^{1/2} \right] \\ &\times \exp \left[\frac{1}{4} \left(\frac{v_f^{(1)}}{\sqrt{2\pi}} \right)^2 (\ln(t/d))^{-1/2} \right] \end{aligned}$$

$$\times \exp \left[-\left(\frac{1}{2} + r \frac{v_f^{(2)}}{\sqrt{2\pi}} \right)^2 (\ln(t/d))^{1/2} \right] \times t^{-1/2} \dots \quad (3.9)$$

It is remarkable that the impurity effect disappears from the most dominant term $\exp[-\frac{1}{3} \ln^{3/2}(t/d)]$; namely neither of $v_f^{(1)}$ and $v_f^{(2)}$ contributes to the dominant term in the FES. This peculiar phenomenon is explained as follows. For the FES correlation function an electron is added to the conduction band at $t = 0$, which then gives rise to the $1/r$ potential scattering to the remaining conduction electrons. This in turn causes the dominant singularity $\exp[-\text{const.} \ln^{3/2}(t/d)]$ in the long-time behavior, hiding the contribution from the core hole created. This conclusion is consistent with the result deduced by Fabrizio et al. for a static hole. [10]

3.2 Crossover to the Wigner crystal: $V(r) = 1/r^\alpha$ case

Now that the critical behavior in the Wigner crystal has been clarified, let us study the TL liquid with $V(r) = 1/(r^2 + d^2)^{\nu+1/2}$ interaction ($\nu > 0$) in order to see the crossover to the Wigner crystal.

3.2.1 $1/r^\alpha$ impurity potential

We first deal with the case of $\tilde{V}(p) = V(p)$; i.e. the impurity potential takes the same form as that for the interaction among conduction electrons. From Eq.(2.3), one can see $V(p) \gg 1$ in the limit $\nu \rightarrow 0$, which results in $\Theta(p) \simeq 1/\sqrt{2}$. Therefore, we end up with $t^{-\alpha(\nu, d)}$ -type *power-law* behavior with the exponents both for OC ($\alpha_{OC}(\nu, d)$) and for FES ($\alpha_{FES}(\nu, d)$), which are obtained as,

$$\alpha_{OC}(\nu, d) = \frac{1}{4} \left(\frac{v_f^{(1)}}{\sqrt{2\pi}} \right)^2 [2F(\nu, d)]^{\frac{1}{2}} + \left(\frac{v_f^{(2)}}{\sqrt{2\pi}} \right)^2 [2F(\nu, d)]^{-\frac{1}{2}} \quad (3.10)$$

$$\begin{aligned} \alpha_{FES}(\nu, d) = & \frac{1}{4} \left(1 + \frac{v_f^{(1)}}{\sqrt{2\pi}} \right)^2 [2F(\nu, d)]^{\frac{1}{2}} \\ & + \left(\frac{1}{2} + r \frac{v_f^{(2)}}{\sqrt{2\pi}} \right)^2 [2F(\nu, d)]^{-\frac{1}{2}} + \frac{1}{2}, \quad (3.11) \end{aligned}$$

where $F(\nu, d)$ is defined in Eq.(2.3). Note that the above formulae for the critical exponents are typical for the TL liquid. Therefore, if the host interaction decays faster than $1/r$ (i.e. $\nu > 0$), the FES correlation is governed by the power-law behavior, as should be the case for the TL liquid. In this case, both of the asymmetric scattering $v_f^{(2)}$ and the symmetric scattering $v_f^{(1)}$ contributes to the dominant power-law behavior. Notice, however, that $F(\nu, d)$ becomes very large, in the limit of $\nu \rightarrow 0$, which is seen from the expanded formula,

$$F(\nu, d) = \frac{1}{2d^\nu} \sum_{n=0}^{\infty} \frac{(2n-1)!!}{(2n)!!(\nu+n)}. \quad (3.12)$$

As a result, the first term with $v_f^{(1)}$ in Eqs. (3.10) and (3.11) becomes more dominant than the second terms with $v_f^{(2)}$. This implies that the symmetric scattering becomes more prominent as the system approaches the Wigner crystal. In other words, the effect of the core-hole motion, which emerges via the asymmetric scattering $v_f^{(2)}$, is dramatically suppressed for the FES in the limit $\nu \rightarrow 0$, though it plays the crucial role in the ordinary TL liquids. [15] Finally at $\nu = 0$, the most dominant term is replaced by $\exp[-\text{const.} \ln^{3/2}(t/d)]$ as shown in (3.7), which is typical for the long-time behavior due to the static (localized) hole.

3.2.2 δ -function impurity potential

Even if the impurity potential is replaced by the one with δ -function type ($\tilde{V}(p) = 1$), we find that the correlation functions show the power-law behavior inherent in TL liquids, whose exponents are given by the Gaussian formula like Eq.(3.10) and Eq.(3.11). Therefore, as far as the system stays in the TL liquid, the FES singularity is governed by the power-law behavior. However, as mentioned in the previous subsection, if we consider the Wigner crystal with $1/r$ interaction, there exist qualitative differences between the $1/r$ impurity potential and δ -function impurity potential.

3.3 Green function

To close this section, we briefly mention the long-distance behavior of the one-particle Green function (GF), which was discussed by Schulz. [9] The long-distance critical behavior of the GF for $\nu > 0$ (TL liquid) is simply obtained by setting $v_f^{(1)} = v_f^{(2)} = 0$ in the FES exponent (3.11),

$$\alpha_G(\nu, d) = \frac{1}{4}[2F(\nu, d)]^{\frac{1}{2}} + \frac{1}{4}[2F(\nu, d)]^{-\frac{1}{2}} + \frac{1}{2} \quad (3.13)$$

This exponent takes a typical form for the GF of the TL liquid,

$$\frac{1}{4} \left(\frac{1}{K_\rho} + K_\rho \right) + \frac{1}{2}, \quad (3.14)$$

with $K_\rho = [2F(\nu, d)]^{-\frac{1}{2}}$. On the other hand, by performing a similar procedure in Eq.(3.7), we obtain the GF for the case of $1/r$ long-range interaction,

$$\begin{aligned} & \langle \psi_r(x) \psi_r^\dagger(0) \rangle \\ &= \exp \left[-\frac{1}{3} (\ln(x/d))^{3/2} \right] \exp \left[-(\ln(x/d))^{1/2} \right] \dots \\ & \times x^{-1/2}, \end{aligned} \quad (3.15)$$

which reproduces the formula obtained by Schulz for the 1D Wigner Crystal. [9]

It is now instructive to discuss the above GF in the viewpoint of a mobile core hole treated here, which allows us to clearly see the origin of the characteristic behavior in the GF for the Wigner crystal. Let us first notice that for $\nu > 0$ the critical exponent (3.14) for

the charge sector has two contributions, the one coming from charged excitations, $1/K_\rho$, and the other from the current excitations, K_ρ . By observing the formula (3.11), we can see that these should be referred to as the contributions from the symmetric and the asymmetric impurity scatterings, respectively, if we regard the added electron in the system as a mobile impurity. As $\nu \rightarrow 0$, $1/K_\rho$ becomes very large, and finally at $\nu = 0$ these two terms are respectively changed to the anomalous critical behaviors $\exp[-\ln^{3/2} x]$ and $\exp[-\ln^{1/2} x]$ inherent in the Wigner crystal. [9] Therefore, we can naturally identify these anomalous contributions with those from the charged (symmetric forward scattering) and the current excitations (asymmetric forward scattering). Based on these correspondences, we can say that the reason why the term $\exp[-\ln^{1/2} x]$ is less dominant than $\exp[-\ln^{3/2} x]$ is due to the fact that the asymmetric forward scattering is to be short-ranged even if the the microscopic interaction is long-ranged, as mentioned in §2.

§4. Summary

We have studied 1D electron systems with $1/r^\alpha$ long-range interaction to see how the TL liquid exhibits a crossover to the Wigner crystal. In particular, we have focused on the FES phenomena with a *mobile* core hole suddenly created for the valence band, which contains the ordinary symmetric forward scattering as well as the asymmetric scattering. It has been shown that, if both of the interactions among the host electrons and the impurity potential is of $1/r^\alpha$ type, the ordinary symmetric scattering becomes more dominant than the asymmetric scattering as the system approaches the Wigner crystal. This is quite contrasted to the ordinary TL liquid for which both of the symmetric and asymmetric potential scatterings play an important role in the FES problem. On the other hand, if the impurity scattering due to the mobile core hole are assumed to be short ranged while the host electrons have the $1/r$ long-range interaction, both of the symmetric and asymmetric scatterings, are entirely suppressed for the FES in the Wigner crystal, being in contrast to the case with long-range impurity potential.

Acknowledgements

We would like to thank T. Fujii for valuable discussions. This work was partly supported by a Grant-in-Aid from the Ministry of Education, Science, Sports and Culture, Japan.

-
- 1) F. D. M. Haldane: J. Phys. C **14**, (1981) 2585; Phys. Rev. Lett. **47**, (1981) 1840.
 - 2) V.J. Emery: in *Highly Conducting One-Dimensional Solids*, edited by J.T. Devreese *et al.* (Plenum, New York, 1979).
 - 3) H. Fukuyama and H. Takayama: *Electronic Properties of Inorganic Quasi-One-Dimensional Compounds*, eds. P. Monceau (D. Reidel, 1985), p.41.

- 4) J. Voit: Rep. Prog. Phys. **58**, (1995) 977.
- 5) A.R. Goñi, A. Pinczuk, J.S. Weiner, J.M. Calleja, B.S. Dennis, L.N. Pfeiffer, and K.W. West: Phys. Rev. Lett. **67**, (1991) 3298.
- 6) R. Egger and A.O. Gogolin: Phys. Rev. Lett. **79**, (1997) 5082.
- 7) C. Kane, L. Balents and M.P.A. Fisher: Phys. Rev. Lett. **79**, (1997) 5086.
- 8) H. Yoshioka and A.A. Odintsov: Phys. Rev. Lett. **82**, (1999) 374.
- 9) H. J. Schulz: Phys. Rev. Lett. **71**, (1993) 1864.
- 10) M. Fabrizio, A. O. Gogolin, and Scheidl: Phys. Rev. Lett. **72**, (1994) 2235.
- 11) H. Maurey and T. Giamarchi: Phys. Rev. B **51**, (1995) 10833.
- 12) R. Egger and H. Grabert: Phys. Rev. Lett. **75**, (1995) 3505.
- 13) H. Otani and T. Ogawa: Phys. Rev. B **54**, (1996) 4540.
- 14) Q. Yuan, H. Chen Y. Zhang and Y. Chen: Phys. Rev. B **58**, (1998) 1084.
- 15) Y. Tsukamoto, T. Fujii and N. Kawakami: Phys. Rev. B **58**, (1998) 3633; *ibid.* Eur. Phys. B **5**, (1998) 479.
- 16) P. L. McEuen, E. B. Foxman, J. Kinaret, U. Meirav, M. A. Kastner, N. S. Wingreen and S. J. Wind: Phys. Rev. **B45** (1992) 11419.
- 17) T. Ogawa, A. Furusaki and N. Nagaosa: Phys. Rev. Lett. **68**, (1992) 3638.
- 18) G.D. Mahan: Phys. Rev. **153**, (1967) 882.
- 19) K.D. Schotte and U. Schotte: Phys. Rev. **182**, (1969) 479.
- 20) K. Ohtaka and Y. Tanabe: Rev. Mod. Phys. **62**, (1990) 929 .
- 21) I. Affleck and A.W.W. Ludwig: J. Phys. A. **27**, (1994) 5375.